

PRODUCT INFORMATION

DEUCE is a selective herbicide for the control of various broadleaf weeds and poa annua in asparagus, garlic, leeks, onions, and potatoes. DEUCE is absorbed through both the leaves and roots of the weeds providing both contact and soil residual activity. Do not apply to light soils, especially in wet conditions, as damage may occur in pre-emergence applications.

WEEDS CONTROLLED:**SUSCEPTIBLE:**

Black Nightshade	Charlock	Chickweed	Fathen	Field Speedwell
Fumitory	Gaiusoga	Groundsel	Henbit	Nettle
Plantains	Annual Poa	Scentless Chamomile	Shepherds Purse	Spurrey
Stinking Mayweed	Twin Cross	White Campion	Wild Radish	Wild Turnip
Willow Weed				

MODERATELY SUSCEPTIBLE:

Combind	Red Root (over 4 leaf)	Sow Thistle	Wire Weed	Seedling Docks
Seedling Californian Thistle				

RESISTANT:

Cleavers	Docks	Giant Buttercup	Rayless Chamomile	most deep rooting weeds
----------	-------	-----------------	-------------------	-------------------------

DIRECTIONS FOR USE:

APPLICATION: Apply through conventional ground equipment ensuring good coverage of the weeds. Half fill the spray tank and, with agitation running, add the required amount of DEUCE. Complete filling of the spray tank with agitation running to ensure complete mixing before spraying.

RECOMMENDATIONS

CROP	TIMING	REMARKS
ASPARAGUS	Seed beds and first year direct seeded	Apply 2 kg per hectare in 340 - 450 litres of water pre-emergence or post emergence when the crop is 10 - 15 cm high. Do not use during the cutting season.
GARLIC	Post planting	Apply 3 kg per hectare before crop emerges.
	Post emergence	Apply 2 kg per hectare once the crop has 3 or more true leaves. Refer onions for split application method. Apply in 340 - 450 litres water per hectare.
ONIONS	Post emergence	Apply 1.5 - 2 kg per hectare in 340 - 450 litres of water/ha when onions have 3 true leaves or more. Re-apply as required.
SPLIT APPLICATIONS		Apply 2 - 3 sprays at rates of 250 - 500 gm/ha in 340 - 450 litres of water at 7 day intervals in combination with other herbicides registered for onions, eg. IOTYRL®91, to improve the spectrum of activity.
LEEKS	Post emergence	Apply 2 kg per hectare once plants become established after transplanting, in 340 - 450 litres of water.
POTATOES	Pre-emergence	Apply 2.5 - 3 kg per hectare in 340 - 450 litres of water before the potatoes emerge.

COMPATIBILITY:

Do not combine with emulsifiable concentrates or adjuvants unless previous experience has proven that the products are compatible and crop safety is not lost.

140mm wide

CROP ROTATION:

Do not plant crops other than those recommended for treatment on this label within 3 months of applying Trump. To minimise residual effects, treated areas should be ploughed and thoroughly cultivated before sowing following crops. liable for a consistent quality of the product but the risk of its storage and use is not borne by us.

SPRAY DRIFT: Do not allow spray drift over aquatic environment. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with any relevant regional air management plans. Aerial applications - to minimise the risk of damage from drift, especially where susceptible vegetation is within 50 metres, use low drift- nozzles which minimise the production of small droplets (<100 microns). Preferably spray in a constant light breeze that is away from sensitive crops.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Elliott Technologies Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

24 Hour emergency response
0800 CHEMCALL (0800 243 622)

© - Registered Trademark of Elliott Technologies Limited
EL144

Batch No.:

DOM:



9 421022 421085

80mm wide

DEUCE™

WDG

A Selective herbicide for broadleaf weed control in asparagus, garlic, leeks, onions and potatoes

Active Ingredient: 700 g/kg methabenzthiazuron in the form of a water dispersible granule.

Read the Label Carefully before use

NET CONTENTS:
1 Kg

Registrant: Elliott Technologies Limited
45 Kitchener Road, Pukekohe
Auckland New Zealand
Telephone: 0800 100 325
www.elliott-technologies.co.nz

Registered pursuant to the ACVM Act 1997, No.4975
HSNO Approval Code H5R000537

HERBICIDE

140mm wide

HARMFUL

Keep out of reach of children

ECOTOXIC
Aquatic Organisms

HSNO Class: 6.4A, 9.1A, 9.2A

Approved Handler Applies

This product must be under the personal control of an approved handler or secured

ENVIRONMENTAL WARNING: VERY TOXIC TO AQUATIC ORGANISMS.

Do not apply to bodies of water. Do not contaminate sewers, drains, dams, creeks or any other waterways with product or empty container. VERY TOXIC IN SOILS - herbicide.

HEALTH WARNING: HARMFUL - may cause eye irritation. Avoid contact with eyes. Wash hands and exposed skin before meals and after work.

PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Burn if circumstances, especially wind direction allow, otherwise bury in landfill. Burning of agricultural containers may require a permit from a Regional Council.

HANDLING PRECAUTIONS: When mixing or applying, wear protective clothing, gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work. Store in original container away from food and feedstuffs. Flush spray equipment thoroughly with clean water after use.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. This product is classified under HSNO as 6.1D and 9.1A, and storage of certain quantities may require bunding and signage. See the HAZNOTE for further information.

When stored appropriately this product should show no significant degradation for 2 years. Contact your supplier for further information about the use of any product that is older than this.

First Aid: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 Poisons (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

165mm deep

Dieline

PMS 485

PMS 369

Black

80mm wide