



# MIZAR®

# GRANUFLO

A Fungicide for the control of Black Spot, Glomerella, and Ripe Spot in Pipfruit and Peach Leaf Curl, and Rust in Stonefruit

**Active Ingredient:** 760 g/kg ziram in the form of a water dispersible granule.

**NET CONTENTS 10 kg**

21039832

Registrant: Elliott Technologies Limited  
45 Kitchener Road, Pukekohe  
New Zealand  
Phone: 0800 100 325



Registered pursuant to ACVM Act 1997, No. P4176  
Approved pursuant to HSNO Act 1996, Approval Code HSR000585

**Fungicide**

® Registered TM of Taminco  
®1 Registered TM of Elliott Technologies Ltd

# HARMFUL

# ECOTOXIC

## KEEP OUT TO REACH OF CHILDREN

## AQUATIC ORGANISMS

**HSNO Class: 6.1D, 6.4A, 6.5B, 6.9B, 9.1A, 9.3C**  
**Approved Handler and Tracking Apply**

**Environmental Warning:** Ecotoxic to aquatic life. Toxic to fish. Do not apply to bodies of water. Do not contaminate sewers, drains, dams, creeks or any other waterways with product or empty container. Harmful to terrestrial vertebrates.

**Health Warning:** Harmful – may be harmful if swallowed, inhaled or absorbed through skin. May cause eye & skin irritation. Wash hands and exposed skin before meals and after work. Believed to cause harm to the liver, central nervous system, and thyroid from repeated oral exposure at high doses.

**Container Disposal:** Triple rinse container and add residue to spray tank. Burn if circumstances, especially wind direction allow, otherwise bury in landfill. Burning of agricultural containers may require a permit from a Regional Council.

**Handling Precautions:** When mixing or applying, wear protective clothing, gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work. Store in original container away from food and feedstuffs.

**First Aid:** If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

**24 Hour Emergency Response 0800 CHEM CALL**  
**0800 236 2255**

### General Information:

Mizar Granuflo is a dry flowable fungicide particularly active against black spot and summer rots, including glomerella and ripe spot in pipfruit and peach leaf curl and rust in stonefruit. Mizar Granuflo provides a protectant coating on plant surfaces preventing disease from establishing. Complete spray coverage is essential.

### Directions for Use:

Mixing: Pour the required amount into the partly filled spray tank with agitation running. Complete filling the spray tank.

**Recommendations:** The dilution rates suggested are for high volume spraying to run-off and are expressed as grams/100 litres of water. Because Mizar Granuflo has a uniform bulk density, it can be poured and measured like a liquid. The volumetric equivalent rates are shown in brackets. For concentrate spraying, adjust dilution rate accordingly. Use the higher rate when conditions favour disease.

| Crop                       | Disease                               | Rate per 100 litres water  | Remarks  |
|----------------------------|---------------------------------------|----------------------------|--|
| Pipfruit                   | Black spot<br>Glomerella<br>Ripe spot | 150 g (285 ml)             | For mid-season protection apply up to three sprays after 3 <sup>rd</sup> cover on a 10 – 14 day schedule                         |
| Stone fruit<br>Pre-blossom | Peach leaf curl                       | 300 – 450 g (570 – 855 ml) | Apply 1-2 sprays pre-blossom. Do not mix with copper fungicides.   |
| Stonefruit<br>Post-blossom | Peach leaf curl, rust                 | 150 – 200 g (285 – 380ml)  | Apply at 10 – 21 day intervals as a preventative spray. Use high rate and short interval during periods of high disease pressure |

### Compatibility:

Compatible with most commonly used pesticides except copper compounds, Bordeaux mixture and lime Sulphur.

**Withholding periods:** Pipfruit – 28 days; Stonefruit – 35 days. Check with export authorities for withholding period on fruit destined for export.

**Storage:** Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. Storage should be in accordance with NZS8409 Management of Agrichemicals. This product is classified under HSNO controls 6.1D and 9.1A, as such storage of certain quantities of this class may require bunding and signage. See the Haznote for further information.

When stored appropriately this product should show no significant degradation for 2 years. Contact your supplier for further information about the use of any product that is older than this.

### Exclusion of Liability: Please Read

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Elliott Technologies Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

**Batch No.**

**DOM**

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.