

KEEP OUT OF REACH OF CHILDREN
Read the Label Carefully before use



SuperSix^{®1}

For the control of powdery mildew on grapes,
tamarillos, peas and vegetables;
rust in peaches, plums and nectarines;
and powdery mildew and rust in
roses and ornamentals

Active Ingredient:
720 g/litre sulphur in the form of a suspension concentrate.

NET CONTENTS:
10 Litres



Registered By:
Elliott Technologies Limited
PO Box 838, Pukekohe, New Zealand.
Phone: 0800 100 325
www.elliott-technologies.co.nz



180mm deep

150mm wide



Dieline



PMS 485



Pantone Purple



Black

HSNO Class: 6.4A, 9.1D

ENVIRONMENTAL WARNING: TOXIC TO FISH.
Avoid contamination of sewers, drains, dams,
creeks or any other waterways with product or
empty container.

**HEALTH WARNING: HARMFUL – MAY CAUSE EYE
IRRITATION.** Avoid contact with eyes.

PRODUCT DISPOSAL: Ensure unwanted product
is used by another operator strictly in accordance
with label directions.

CONTAINER DISPOSAL: Triple rinse container and
add residue to spray tank. Burn if circumstances,
especially wind direction allow, otherwise bury in
landfill. Burning of agricultural containers may
require a permit from a Regional Council.

CLEANING: Flush spray equipment thoroughly
with clean water after use.

HANDLING PRECAUTIONS: When mixing or
applying, wear protective clothing, gloves and
goggles. Do not eat drink or smoke while using.
Wash hands and face before meals and after work.
Wash protective clothing after work. Flush spray
equipment thoroughly with clean water after use.

STORAGE: Store in original container tightly
closed and in a locked, dry, cool, well ventilated
area, away from feed, seeds and foodstuffs.
Storage must be in accordance with NZS8409
Management of Agricultural Chemicals. See the HAZNOTE
for further information.

When stored appropriately this product should
show no significant degradation for 2 years.
Contact your supplier for further information about
the use of any product that is older than this.

FIRST AID: If swallowed do NOT induce vomiting.
For advice, contact the **NATIONAL POISONS CENTRE
0800 POISON (0800 764 766)** or a doctor
immediately. If skin or hair contact occurs, remove
contaminated clothing and flush skin and hair with
running water.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users
of the goods accept that neither the manufacturer,
Elliott Technologies Limited as distributor, nor any
other distributor have any liability or responsibility
whatsoever for any loss, damage or injury whether
in contract or tort, whether direct, indirect or
consequential howsoever arising in connection with
the supply of these goods.

Batch No.

DOM

PRODUCT INFORMATION

SUPER SIX is a flowable sulphur formulation which combines ease of use with optimum
particle size and excellent adhesion. **SUPER SIX** is a contact fungicide with a broad spectrum
of activity and a good effect against Erinose mite in Grapes. When used as directed **SUPER SIX**
assists where sulphur is deficient.

DIRECTIONS FOR USE

MIXING: Pour the required amount into the partly filled spray tank with agitation running,
complete filling of spray tank. Always shake the container before use.

RECOMMENDATIONS:

The dilution rates suggested are for high volume spraying to run-off and are expressed as
mls/100 litres of water. For concentrate spraying adjust dilution rate accordingly. Use the
higher rate when conditions favour disease.

CROP	DISEASE	RATE/100L	REMARKS
Grapes	Powdery Mildew Erinose Mite	165 – 330 ml	Apply at 10-14 day intervals from bud swell To six weeks before harvest
Peaches Plums Nectarines	Rust	110 – 220 ml	Apply at 3-weekly intervals from fruit set until harvest. Do not apply to Apricots or mature Nectarines.
Apples	Powdery Mildew	110 – 220 ml	Apply from bud movement until harvest. NOTE: (1) Do not mix with Captan on Sulphur sensitive varieties such as Cox's Orange Pippin, Kidd's Orange or Red Delicious (2) It is not advisable to use Super Six post blossom unless one spray has been used pre-blossom. (3) Fruit finish may be affected on some varieties. (4) Do not use above 25°C (5) Do not use on Pears.
Tamarillos	Powdery Mildew	110 – 220 ml	Apply at 10 – 14 day intervals throughout the season
Peas	Powdery Mildew	110 – 220 ml	Apply when conditions favour disease and repeat as necessary.
Vegetables	Powdery Mildew	110 – 220 ml	Apply on a 14 day intervals as required. Do not apply to Rock Melons or Cucumbers
Roses and Ornamentals	Powdery Mildew	220 – 330 ml	Apply on a 14 day schedule. Do not mix with maldison.

COMPATIBILITY: Do not combine with oils or use within 21 days of an oil spray. Super Six can
be applied with liquid fertilisers but caution should be exercised if mixing with previously
unried emulsifiable concentrates.

WITHHOLDING PERIOD: Nil - all crops.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

® is a registered trademark of Elliott Technologies Ltd

®1 Super Six is a registered TM of Griffin Corporation USA

Registered pursuant to the ACVM Act 1997, No. P3979

Approved pursuant to HSN Act 1996, Approval Code HSR000666

24 Hour Emergency Response
0800 CHEM CALL
0800 243 622



EL216

9 421022 420996

180mm deep

150mm wide



Dieline



Black